



Hermite, Higher order Hermite, Laguerre type polynomials and Burgers like equations

Giuseppe Dattoli, Roberto Garra*, Silvia Licciardi

ENEÀ - Frascati Research Center, Via Enrico Fermi 45, 00044, Frascati, Rome, Italy

Università Telematica Internazionale, Section of Mathematics, Italy

University of Palermo, Department of Engineering, viale delle Scienze, Palermo, Italy

ARTICLE INFO

MSC:

58J35

33C45

35K05

42C05

Keywords:

Burgers-type differential equations

Hermite polynomials

Laguerre polynomials

ABSTRACT

The multivariable version of ordinary and generalized Hermite polynomials are the natural solutions of the classical heat equation and of its higher order versions. We derive the associated Burgers equations and show that analogous non-linear partial differential equations can be derived for Laguerre polynomials. The starting point of this extension is the Laguerre diffusive equation, whose nonlinear extension reveals interesting implications involving families of mixed polynomials. In this way we have a general scheme to obtain new exact explicit solutions for nonlinear PDEs by using Laguerre, Hermite and other families of polynomials of Appèl and non-Appèl type.

1. Introduction and preliminaries

The theory of special functions and polynomials has been the subject of intensive programs of research during the years. Most of these studies have been directed towards the framing of the relevant theory within a unifying context. Which from time to time has been identified with the theory of differential equations [1], the group theoretic point of view [2], umbral calculus [3] and symbolic formalism [4], just to quote a few of them. An important output of these studies has been the formalization of the theoretical foundations of the relevant properties in terms of differential equations, be they ordinary or partial. Multi-variable special functions are the natural solutions of well-known equations of mathematical physics and the possibility of extending these equations to non-linear forms is the subject of the present study.

The two variable Hermite polynomials are defined by [5]

$$H_n(x, y) = n! \sum_{r=0}^{\lfloor \frac{n}{2} \rfloor} \frac{y^r x^{n-2r}}{(n-2r)! r!}. \quad (1.1)$$

and satisfy the parabolic equation

$$\frac{\partial Z}{\partial y} = \frac{\partial^2 Z}{\partial x^2} \quad (1.2)$$

under the condition $Z(x, 0) = x^n$. Albeit two variable Hermite are well known in the mathematical literature and trace back to Hermite himself [6], Appèl and Kampè dè Fèrèrè [7], they received a wider diffusion when they have been exploited as auxiliary

* Corresponding author at: Università Telematica Internazionale, Section of Mathematics, Italy.

E-mail address: roberto.garra@uninettunouniversity.net (R. Garra).

solutions of the heat equation, under the name of heat polynomials, for more specific details see Ref. [8]. A further element of interest is due to the fact that they can be associated with non-linear differential equations realizing the so called Burgers hierarchy [9–11].

In this paper, we show the applications of Hermite and Laguerre polynomials to solve nonlinear Burgers-type equations by using the Hopf-Cole transform. We also consider nonlinear equations involving Laguerre derivatives, showing interesting possibilities of applications of families of mixed polynomials. We finally provide further examples and considerations in order to highlight the potential impact of the theory of special polynomials to constructs solutions for nonlinear PDEs.

2. Applications of Hermite polynomials to solve nonlinear PDEs

In this section we show that it is possible to construct particular interesting solutions for a wide class of nonlinear partial differential equations by using Hermite and Hopf-Cole transformation (HCT). We introduce the problem through the following example.

Example 1. We recall that, by applying the HCT [11], we obtain

$$u(x, y) = \frac{\partial Z}{\partial x} \Big/ Z = \frac{\partial}{\partial x} \ln Z, \tag{2.1}$$

with $Z(x, y)$ given by (1.1) satisfying Eq. (1.2). We can easily show that the function $u(x, y)$ satisfies a non-linear equation presenting simultaneously non-linear advective and diffusive terms (the so called Burgers equation). Keeping the partial derivative of with respect to the y -variable, we obtain

$$\frac{\partial u}{\partial y} = \frac{\left(\frac{\partial^2 Z}{\partial y \partial x} \right) Z - \frac{\partial Z}{\partial y} \frac{\partial Z}{\partial x}}{Z^2}. \tag{2.2}$$

The use of Eq. (1.2) yields

$$\frac{\partial u}{\partial y} = \frac{\left(\frac{\partial^3 Z}{\partial x^3} \right) Z - \frac{\partial^2 Z}{\partial x^2} \frac{\partial Z}{\partial x}}{Z^2}. \tag{2.3}$$

Then, according to Eq. (2.1), we find that the function u satisfies the equation

$$\frac{\partial u}{\partial y} = \frac{\partial^2 u}{\partial x^2} + 2u \frac{\partial u}{\partial x}. \tag{2.4}$$

This means that the Burgers Eq. (2.4) admits a solution of the form (see also [10])

$$u(x, y) = \frac{nH_{n-1}(x, y)}{H_n(x, y)}. \tag{2.5}$$

This result is a particular case, to be framed within a more general context. For example, the third order heat equation

$$\frac{\partial \Psi}{\partial t} = \frac{\partial^3 \Psi}{\partial x^3}, \tag{2.6}$$

is associated with a non-linear equation, generalizing the already mentioned Burgers. The use of the same transformation given in Eq. (2.1) and extension of the procedure outlined afterwards allows to prove that the function u satisfies the non-linear form [10]

$$\frac{\partial u}{\partial y} = \frac{\partial u^3}{\partial x} + 3 \left(\frac{\partial u}{\partial x} \right)^2 + 3u \frac{\partial^2 u}{\partial x^2} + \frac{\partial^3 u}{\partial x^3} \tag{2.7}$$

which, in a more compact form, reads

$$\frac{\partial u}{\partial y} = \frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} + u \right)^2 u. \tag{2.8}$$

We recall that the nonlinear Eq. (2.7) is known in the literature as the Tasso–Sharma–Olver equation. This equation is a relevant example of odd members of the Burgers hierarchy that can be transformed in a linear equation involving highest derivatives. It was firstly studied in [12,13]. There is a wide literature about this nonlinear PDE, we refer for example to [14] and the references therein.

Observation 1. The last Eqs. (2.7) and (2.8) are obtained by proceeding on the basis of the paradigm reported below:

(i) introduce the auxiliary function defined as in Eq. (2.1), namely

$$u(x, y) = \frac{\partial \Psi}{\partial x} \Big/ \Psi, \tag{2.9}$$

(ii) keep the derivative of both sides of Eq. (2.9) with respect to and obtain

$$\frac{\partial u}{\partial y} = \frac{\left(\frac{\partial^2 \Psi}{\partial y \partial x} \right) \Psi - \frac{\partial \Psi}{\partial y} \frac{\partial \Psi}{\partial x}}{\Psi^2} = \frac{\left(\frac{\partial^4 \Psi}{\partial x^4} \right) \Psi - \frac{\partial^3 \Psi}{\partial x^3} \frac{\partial \Psi}{\partial x}}{\Psi^2}, \tag{2.10}$$

(iii) use the identity

$$\frac{\partial \Psi}{\partial x} = u\Psi. \tag{2.11}$$

In Eq. (2.10), eventually ending up with

$$\begin{aligned} \frac{\partial u}{\partial y} &= \frac{\frac{\partial^3(u\Psi)}{\partial x^3} - u \frac{\partial^3\Psi}{\partial x^3}}{\Psi} = \frac{\Psi \frac{\partial^3 u}{\partial x^3} + 3 \frac{\partial \Psi}{\partial x} \frac{\partial^2 u}{\partial x^2} + 3 \frac{\partial u}{\partial x} \frac{\partial^2 \Psi}{\partial x^2}}{\Psi} \\ &= \frac{\partial^3 u}{\partial x^3} + 3u \frac{\partial^2 u}{\partial x^2} + \frac{3}{\Psi} \frac{\partial u}{\partial x} \frac{\partial(u\Psi)}{\partial x} \end{aligned} \tag{2.12}$$

Expanding the last term on the rhs of the last line in (2.12), we eventually end up with Eq. (2.7).

As is well known the third order Hermite polynomials [5]

$$H_n^{(3)}(x, y) = n! \sum_{r=0}^{\lfloor \frac{n}{3} \rfloor} \frac{y^r x^{n-3r}}{(n-3r)!r!} \tag{2.13}$$

satisfies Eq. (2.6) and

$$\frac{\partial H_n^{(3)}(x, y)}{\partial x} = nH_{n-1}^{(3)}(x, y), \tag{2.14}$$

therefore, we have that a particular solution for the nonlinear Eq. (2.8) is given by

$$u(x, y) = \frac{nH_{n-1}^{(3)}(x, y)}{H_n^{(3)}(x, y)}. \tag{2.15}$$

This scheme can be used to construct exact particular solutions for higher order nonlinear PDEs.

The extension of the method to higher order ‘‘heat’’ equations is fairly straightforward. We find indeed that the higher order Burgers-type equations

$$\frac{\partial u}{\partial y} = \frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} + u \right)^{m-1} u \tag{2.16}$$

can be linearized by means of the transform (2.9) to the higher order diffusive equation

$$\frac{\partial \Psi}{\partial y} = \frac{\partial^m \Psi}{\partial x^m}, \quad m > 3. \tag{2.17}$$

It is well-known that higher order diffusive equations of this type admits polynomial solutions of the form

$$H_n^{(m)}(x, y) = n! \sum_{r=0}^{\lfloor \frac{n}{m} \rfloor} \frac{y^r x^{n-mr}}{(n-mr)!r!}. \tag{2.18}$$

Therefore, by using the HCt, it is possible to solve higher order Burger-type Eqs. (2.16) in a similar way.

In this article we pursue an investigation started in Ref. [15], where the authors have studied extensions of the Burgers equations including non-standard derivative forms. Our intention is that of exploring this subject in a wider context, by an appropriate discussion involving not only Hermite but also Laguerre and other forms of hybrid polynomials.

To conclude this section, we observe another interesting application of this method in order to obtain solutions based on Hermite polynomials for a more general class of Burgers equations.

Example 2. In the recent paper [16], the author considered the following Burgers-type equation with a variable coefficient

$$\frac{\partial u}{\partial t} + \frac{2}{F(y, t)} u \frac{\partial u}{\partial x} = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2}. \tag{2.19}$$

Considering the following Hopf-Cole transformation

$$u(x, y, t) = -F(y, t) \frac{\partial}{\partial x} \ln(\varphi(x, t)), \tag{2.20}$$

we have that the function $F(y, t)$ and $\varphi(x, t)$ should solve diffusive equations, respectively

$$\frac{\partial \varphi}{\partial t} = \frac{\partial^2 \varphi}{\partial x^2} \tag{2.21}$$

and

$$\frac{\partial F}{\partial t} = \frac{\partial^2 F}{\partial y^2}. \tag{2.22}$$

Therefore, in view of the connection with the Hermite polynomials, we can construct a particular interesting solution also for the problem of the form

$$u(x, y, t) = -H_n(y, t) \frac{nH_{n-1}(x, t)}{H_n(x, t)}. \tag{2.23}$$

In this section we have fixed the essential tools of the formalism, we will employ these methods in the forthcoming sections, aimed at combining HCT, properties of special polynomials and operational methods to derive various forms of non-linear equations, some of which not previously known.

3. Burgers equations and special polynomials

The polynomials defined in (2.13) are sometimes referred as lacunary third order Hermite, their complete version is provided by

$$H_n^{(3)}(x_1, x_2, x_3) = n! \sum_{r=0}^{\lfloor \frac{n}{3} \rfloor} \frac{H_{n-3r}(x_1, x_2) x_3^r}{(n-3r)! r!} \tag{3.1}$$

satisfying the recurrences

$$\frac{\partial}{\partial x_1} H_n^{(3)}(x_1, x_2, x_3) = n H_{n-1}^{(3)}(x_1, x_2, x_3) \tag{3.2}$$

and

$$\frac{\partial}{\partial x_2} H_n^{(3)}(x_1, x_2, x_3) = \frac{\partial^2}{\partial x_1^2} H_n^{(3)}(x_1, x_2, x_3). \tag{3.3}$$

Moreover,

$$H_n^{(3)}(x_1, 0, x_3) = H_n^{(3)}(x_1, x_3), \quad H_n^{(3)}(x_1, x_2, 0) = H_n(x_1, x_2). \tag{3.4}$$

According to the previously developed discussion, two different Burgers can be associated with the last two heat type equations.

Proposition 3.1. $\forall x_1, x_2, x_3 \in \mathbb{R}, \forall n \in \mathbb{N}$, the following PDEs hold

$$\frac{\partial u_n}{\partial x_3} = \frac{\partial}{\partial x_1} \left(\frac{\partial}{\partial x_1} + u_n \right)^2 u_n \tag{3.5}$$

$$\frac{\partial u_n}{\partial x_2} = \frac{\partial}{\partial x_1} \left(\frac{\partial}{\partial x_1} + u_n \right) u_n. \tag{3.6}$$

Proof. Eqs. (3.5) and (3.6) are easily proved by setting

$$u_n(x_1, x_2, x_3) = \frac{n H_{n-1}^{(3)}(x_1, x_2, x_3)}{H_n^{(3)}(x_1, x_2, x_3)}. \quad \square \tag{3.7}$$

Corollary 3.2. It is furthermore evident that in the case of the nonlacunary m th order Hermite, the following ‘‘hierarchical’’ system of Burgers equation holds

$$\frac{\partial u_n}{\partial x_k} = \frac{\partial}{\partial x_1} \left(\frac{\partial}{\partial x_1} + u_n \right)^{k-1} u_n, \quad 1 < k \leq m, \tag{3.8}$$

that admits the solution

$$u_n(x_1, \dots, x_m) = \frac{n H_{n-1}^{(m)}(x_1, \dots, x_m)}{H_n^{(m)}(x_1, \dots, x_m)}. \tag{3.9}$$

The examples, we have considered so far, deal with polynomials belonging to the Hermite family. The wealth of the existing polynomials can be exploited as a benchmark for the associated Burgers like equations. To this aim we consider the two variable Laguerre polynomials.

We recall that the two-variable Laguerre polynomials

$$L_n(x, t) = n! \sum_{r=0}^n \frac{t^{n-r} x^r}{(n-r)! (r!)^2} \tag{3.10}$$

solves the Laguerre diffusive equation [17,18]

$$\frac{\partial F}{\partial t} = \left(\frac{\partial}{\partial x} x \frac{\partial}{\partial x} \right) F. \tag{3.11}$$

Observation 2. In the recent paper [15], the authors have obtained the related Burgers-type equation by using a scheme similar to the one used in the previous section. Indeed, by using the HCT

$$u = \frac{\partial_x F}{F} \tag{3.12}$$

after keeping the (partial) time derivative of both sides of (3.12) with respect to the variable t , we obtain

$$\frac{\partial u}{\partial t} = \frac{\left(\frac{\partial}{\partial t} \frac{\partial F}{\partial x}\right) F - \frac{\partial F}{\partial x} \frac{\partial F}{\partial t}}{F^2}. \tag{3.13}$$

Being t and x independent variables, the associated derivatives commute, so that, also on account of Eq. (3.11), we find

$$\frac{\partial}{\partial t} \frac{\partial F}{\partial x} = \frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} x \frac{\partial F}{\partial x}\right). \tag{3.14}$$

Thus, going back to (3.13) and by using the fact that $\partial_x F = uF$, we obtain

$$\frac{\partial u}{\partial t} = \frac{1}{F} \left(\frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} (xuF) - u \frac{\partial}{\partial x} x \frac{\partial F}{\partial x}\right)\right). \tag{3.15}$$

Working out the derivatives we eventually end up with

$$\frac{\partial u}{\partial t} = \frac{\partial}{\partial x} x \frac{\partial u}{\partial x} + \frac{\partial u}{\partial x} + \left(1 + x \frac{\partial}{\partial x}\right) u^2 \tag{3.16}$$

which is a non-trivial non-linear reaction–diffusion type equation with known solution, once the explicit solution of the linear Laguerre-type Eq. (3.11) is given. This means that we can obtain interesting exact solutions for the non-trivial nonlinear Eq. (3.16) by means of the Laguerre polynomials. A solution for (3.16) will be given by

$$u_n(x, t) = \frac{\partial_x L_n(x, t)}{L_n(x, t)} \tag{3.17}$$

Observe that another interesting generalization of the Burgers equation is given by

$$\frac{\partial u}{\partial t} = \frac{\partial}{\partial x} x \frac{\partial u}{\partial x} + x \left(\frac{\partial u}{\partial x}\right)^2, \tag{3.18}$$

which can be linearized and reduced to Eq. (3.11) by means of the transformation $u = \ln(h(x, t))$. Also in this case we can construct an explicit solution involving Laguerre polynomials to (3.18) as $u(x, t) = \ln(L_n(x, t))$.

Further examples of special polynomials exhibiting PDE generalizing those presented before, eventually linked to Burgers counterparts, are discussed below.

Example 3. A family of polynomials of “hybrid” type is provided by (see [19])

$${}_2\mathcal{L}_n(x, y) = n! \sum_{r=0}^{\lfloor \frac{n}{2} \rfloor} \frac{y^r x^{n-2r}}{(n-2r)! r!^2}, \tag{3.19}$$

which has both Hermite and Laguerre “imprinting”, for this reasons they satisfy the PDE

$$\frac{\partial}{\partial y} y \frac{\partial}{\partial y} \Psi(x, y) = \frac{\partial^2}{\partial x^2} \Psi(x, y), \tag{3.20}$$

under the condition $\Psi(x, 0) = x^n$. The previous equation deserves a comment. If it is indeed viewed as an initial value problem, its formal solution can be cast in the form

$$\Psi(x, y) = C_0 \left(y \frac{\partial^2}{\partial x^2}\right) x^n, \tag{3.21}$$

where

$$C_0(z) = \sum_{r=0}^{\infty} \frac{z^r}{r!^2} \tag{3.22}$$

is a Bessel like function, satisfying the eigenvalue problem

$$\frac{d}{dz} z \frac{d}{dz} C_0(\lambda z) = \lambda C_0(\lambda z). \tag{3.23}$$

The search for the Burgers like form associated with polynomials of the type (3.19) can be accomplished by the use of the same means exploited for the case of non-hybrid families. The use of the transformation in (3.20)

$$u(x, y) = \ln(\Psi(x, y)) \tag{3.24}$$

leads e.g. to

$$\frac{\partial}{\partial y} y \frac{\partial u}{\partial y} + y \left(\frac{\partial u}{\partial y}\right)^2 = \frac{\partial^2 u}{\partial x^2} + \left(\frac{\partial u}{\partial x}\right)^2. \tag{3.25}$$

In this section we have seen how different forms of Burgers equations can be obtained by handling the PDE satisfied by various families of special polynomials, further comments will be given in the forthcoming concluding section.

4. Final comments

In this article, we have touched different aspects regarding the impact of non-linear PDE on the theory of special polynomials. We have proved that the wealth of the relevant properties reflects itself on a comparably rich plethora of Burgers type equations linearized after a CHT involving special polynomials. In order to provide a further flexibility of the method we consider the further example below.

Example 4. We consider the PDE

$$\begin{cases} \frac{\partial}{\partial y} \Psi(x, y) = \left(\alpha \frac{\partial}{\partial x} + \beta \frac{\partial^2}{\partial x^2} + \gamma \frac{\partial^3}{\partial x^3} \right) \Psi(x, y) \\ \Psi(x, 0) = f(x), \end{cases} \tag{4.1}$$

whose formal solution writes

$$\Psi(x, y) = \exp \left\{ y \left(\alpha \frac{\partial}{\partial x} + \beta \frac{\partial^2}{\partial x^2} + \gamma \frac{\partial^3}{\partial x^3} \right) \right\} f(x). \tag{4.2}$$

Taking into account the generating function of the Higher order Hermite polynomials, that is [5]

$$e^{x_1 t + x_2 t^2 + x_3 t^3} = \sum_{n=0}^{\infty} \frac{t^n}{n!} H_n(x_1, x_2, x_3), \tag{4.3}$$

the explicit solution of Eq. (4.1) reads

$$\Psi(x, y) = \sum_{n=0}^{\infty} \frac{1}{n!} H_n(\alpha y, \beta y, \gamma y) \frac{\partial^n}{\partial x^n} f(x) \tag{4.4}$$

which for $f(x) = x^n$ reduces eventually to [5]

$$\Psi(x, y) = H_n(x + \alpha y, \beta y, \gamma y). \tag{4.5}$$

Accordingly, the function

$$u_n(x, y) = \frac{n H_{n-1}(x + \alpha y, \beta y, \gamma y)}{H_n(x + \alpha y, \beta y, \gamma y)} \tag{4.6}$$

is one of the solutions of the non-linear Burgers associated to the Eq. (4.1)

$$\frac{\partial u_n}{\partial y} = \frac{\partial}{\partial x} \left[\alpha + \beta \left(\frac{\partial}{\partial x} + u_n \right) + \gamma \left(\frac{\partial}{\partial x} + u_n \right)^2 \right] u_n, \tag{4.7}$$

which is also a consequence of the previously quoted ‘‘hierarchical’’ properties.

Before closing the article, we would like to comment on the role of the functions

$$\Phi_n^{(m)}(x, y) = \frac{n H_{n-1}^{(m)}(x, y)}{H_n^{(m)}(x, y)} \tag{4.8}$$

which play a role less trivial than it might appear at first glance. They represent indeed a general form of rational solutions of Burgers like equations. The interest for these form of solutions is motivated by the fact that the non-linear equations we have dealt with are simple analogue of the Navier–Stokes equations, and can be exploited as test for the relevant numerical solutions, see e.g. [20]. Even though the problem deserves a more appropriate treatment, we will afford in a forthcoming paper, we note that, for example

$$\begin{aligned} \Phi_2^{(2)}(x, y) &= \frac{2x}{2y + x^2}, \\ \Phi_4^{(2)}(x, y) &= \frac{4x^3 + 24xy}{x^4 + 12x^2y + 12y^2}, \\ \Phi_4^{(4)}(x, y) &= \frac{6y + x^3}{6xy + \frac{x^4}{4}}, \\ \Phi_9^{(4)}(x, y) &= \frac{9.072 \cdot 10^4 x^2 y^2 + 3.024 \cdot 10^3 x^5 + 9x^8}{6.048 \cdot 10^4 y^3 + 3.024 \cdot 10^4 x^3 y^2 + 504x^6 + x^9} \end{aligned}$$

are rational solutions of the Burgers Eq. (2.4) which can be compared with analogous forms, obtained e.g. in [11,21–23]. The behavior of these functions is given in Fig. 1 where we have reported the plots of $\Phi_n^{(m)}(x, \cdot)$ vs. x , at fixed y for different values of the index n and order m . The relevant behavior is that exhibited by the solutions reported in Ref. [21], where the problem of deriving new analytical solutions of rational type has been addressed. A more exhaustive 3-D view is given in Fig. 2.




Fig. 1. Comparison of $\Phi_n^{(m)}(x, y)$ vs. x for fixed y values and different n and m values.




Fig. 2. Eq. (4.8) for different n and m values.

Remark 4.1. The same considerations apply to the case of the two variable Laguerre. Examples of rational solutions of the Burgers type associated with the Laguerre diffusion equation (given in Eq. (3.17)) are listed below for $n = 3$ and $n = 7$

$$u_3(x, y) = \frac{\frac{x^2}{12} + \frac{xy}{2} + \frac{y^2}{2}}{\frac{x^3}{36} + \frac{x^2y}{4} + \frac{xy^2}{2} + \frac{y^3}{6}} \tag{4.9}$$

$$u_7(x, y) = \frac{\frac{x^6}{3628800} + \frac{x^5y}{86400} + \frac{x^4y^2}{5760} + \frac{x^3y^3}{864} + \frac{x^2y^4}{288} + \frac{xy^5}{240} + \frac{y^6}{720}}{\frac{x^7}{25401600} + \frac{x^6y}{518400} + \frac{x^5y^2}{28800} + \frac{x^4y^3}{3456} + \frac{x^3y^4}{864} + \frac{x^2y^5}{480} + \frac{xy^6}{720} + \frac{y^7}{5040}} \tag{4.10}$$

The relevant plots are given in Fig. 3. The structure is not dissimilar from that exhibited by the Hermite counterparts. More appropriate comments will be provided elsewhere.




Fig. 3. Eqs. (4.9)–(4.10) vs. x for fixed y values.

We have so far discussed the use of Hermite polynomials to construct exact solutions of Burgers equations. This point of view can be reversed. The wealth of the properties of the non-linear transformation we have dealt with can be exploited to obtain previously unknown properties between Hermite polynomials, like e.g.

$$\left(\frac{\partial}{\partial x} + F_n(x, y)\right) F_n(x, y) = S_n(x, y), \quad (4.11)$$

where

$$F_n(x, y) = \frac{(n-1)H_{n-2}(x, y)}{H_{n-1}(x, y)}$$

and

$$S_n(x, y) = \frac{(n-1)(n-2)H_{n-3}(x, y)}{H_{n-1}(x, y)},$$

which is a particular case of more general identities obtainable from the handling of the Burgers auxiliary functions. These problems and the construction of new exact solutions of the Burgers equation in terms of Hermite polynomials will be carefully studied in a forthcoming dedicated paper.

Data availability

No data was used for the research described in the article.

Acknowledgments

Dr. Silvia Licciardi was supported by the following project. Funder: Project funded under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.3 - Call for tender No. 1561 of 11.10.2022 of Ministero dell'Università e della Ricerca (MUR); funded by the European Union – NextGenerationEU. Award Number: Project code PE0000021, Concession Decree No. 1561 of 11.10.2022 adop- ted by Ministero dell'Università e della Ricerca (MUR), CUP - Project title “Network 4 Energy Sustainable Transition – NEST.

References

- [1] J.B. Seaborn, Differential equations and special functions, in: Hypergeometric Functions and their Applications, in: Texts in Applied Mathematics, vol. 8, Springer, New York, NY, http://dx.doi.org/10.1007/978-1-4757-5443-8_2.
- [2] J.D. Talman, Special functions: A group theoretic approach, in: Based on Lectures and with an Introduction By E. P. Wigner, in: Mathematical Physics Monograph Series, W. A. Benjamin Incorporated, New York, NY, USA, Amsterdam, The Netherlands, 1968.
- [3] S. Roman, The umbral calculus, Pure and Applied Mathematics, vol. 111, Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1984.
- [4] S. Licciardi, G. G. Dattoli, Guide to the Umbral Calculus, a Different Mathematical Language, World Scientific, 2022.
- [5] D. Babusci, G. Dattoli, S. Licciardi, E. Sabia, Mathematical Methods for Physicists, World Scientific, 2019.
- [6] Ch. Hermite, Sur un nouveau developpement en séries de fonction, CRAS 5 (1864) 93.
- [7] P. Appell, J. Kampé de Fériét, Fonctions Hypergéométriques et Hypersphériques Polynome d'Hermite, Gauthier-Villars, Paris, 1926.
- [8] D.V. Widder, The Heat Equation, vol. 67, Academic Press, 1976.
- [9] X. Cao, C. Xu, A Bäcklund Transformation for the Burgers Hierarchy, in: Abstract and Applied Analysis, vol. 2010, Hindawi, 2010, January.
- [10] G. Dattoli, D. and Levi, Heat polynomials, umbral correspondence and Burgers equations, 2006, arXiv preprint [nlin/0610076](https://arxiv.org/abs/nlin/0610076).
- [11] N.A. Kudryashov, Generalized Hermite polynomials for the Burgers hierarchy and point vortices, Chaos Solitons Fractals 151 (2021) 111256.
- [12] P.J. Olver, Evolution equation possessing infinite many symmetries, J. Math. Phys. 18 (6) (1977) 1212–1215.
- [13] A.S. Sharma, H. Tasso, Connection between wave envelope and explicit solution of a nonlinear dispersive equation, Report IPP6/158 Ber, MPI für Plasmaphysik (Garching) (1977) 1–10.
- [14] Y. Shang, Y. Huang, W. and Yuan, Bäcklund transformations and abundant exact explicit solutions of the Sharma–Tasso–Olver equation, Appl. Math. Comput. 217 (17) (2011) 7172–7183.
- [15] G. Dattoli, R. Garra, A note on differential equations of logistic type, 2023, arXiv preprint [arXiv:2308.06135](https://arxiv.org/abs/2308.06135).
- [16] B.M. Vaganan, Cole-hopf transformations for higher dimensional Burgers equations with variable coefficients, Stud. Appl. Math. 129 (3) (2012) 300–308.
- [17] G. Dattoli, A. Torre, Operational methods and two-variable laguerre polynomials, Atti Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur. 132 (1998) 1–7.

- [18] G. Dattoli, S. Licciardi, R.M. Pidotella, Theory of generalized trigonometric functions: From Laguerre to Airy forms, *J. Math. Anal. Appl.* 468 (1) (2018) 103–115.
- [19] G. Dattoli, S. Lorenzutta, P.E. Ricci, C. Cesarano, On a family of hybrid polynomials, *Integral Transforms Spec. Funct.* 15 (6) (2004) 485–490.
- [20] D.S. Guzhev, N.N. Kalitkin, Burgers equation is a test for numerical methods, *Matematicheskoe Modelirovanie* 7 (4) (1995) 99–127.
- [21] V.I. Avrutskiy, V.P. Krainov, Rational solutions of $(1+1)$ -dimensional Burgers equation and their asymptotic, 2019, arXiv preprint [arXiv:1910.05488](https://arxiv.org/abs/1910.05488).
- [22] N.A. Kudryashov, D.I. Sinelshchikov, Exact solutions of equations for the Burgers hierarchy, *Appl. Math. Comput.* 215 (3) (2009) 1293–1300.
- [23] M. Zuparic, K. Hoek, Green's functions and the Cauchy problem of the Burgers hierarchy and forced Burgers equation, *Commun. Nonlinear Sci. Numer. Simul.* 73 (2019) 275–290.